

# Fuzzy Neutrosophic Weakly-Generalized Closed Sets in Fuzzy Neutrosophic Topological Spaces



Fatimah M. Mohammed<sup>1,\*</sup> & Anas A. Hijab<sup>1</sup> and Shaymaa F. Matar<sup>1</sup>

<sup>1</sup>Department of Mathematics, College of Education for Pure Sciences,  
Tikrit University, Tikrit, IRAQ

## ARTICLE INFO

Received: 29 / 7 /2018  
Accepted: 27 / 9 /2018  
Available online: 3/1/2019  
DOI: [10.37652/juaps.2022.171813](https://doi.org/10.37652/juaps.2022.171813)

### Keywords:

Fuzzy Neutrosophic set.  
Fuzzy Neutrosophic Topology.  
Fuzzy Neutrosophic Weakly-  
Generalized closed sets.

## ABSTRACT

In this paper, we will define a new class of sets, called fuzzy neutrosophic weakly- generalized closed sets, then we proved some theorems related to this definition. After that, we studied some relations between fuzzy neutrosophic weakly-generalized closed sets and fuzzy neutrosophic  $\alpha$  closed sets, fuzzy neutrosophic closed sets, fuzzy neutrosophic regular closed sets, fuzzy neutrosophic pre closed sets and fuzzy neutrosophic semi closed sets.

## Introduction:

The first use of the concept of fuzzy sets was introduced by Zadeh in 1965 [1]. After that, the fuzzy set theory extension by many researchers. Intuitionistic fuzzy sets (IFS) was one of the extension sets and defined by K. Atanassov in 1983 [2, 3, 4], when fuzzy set gives the degree of membership of an element in the sets, whenever intuitionistic fuzzy sets give a degree of membership and a degree of non-membership. After that, several researches were conducted on the generalizations of the notion of intuitionistic fuzzy sets, one of them was Floretin Smarandache in 2010 [5] when he developed another membership in addition to the two memberships which were defined in intuitionistic fuzzy sets and called it neutrosophic set.

The term of neutrosophic sets was defined with membership, non-membership and indeterminacy degrees. In the last year, (2017) Veereswari [9] introduced fuzzy neutrosophic topological spaces. This concept is the solution and representation of the problems with various fields.

In this paper, We introduced define a new class of sets via fuzzy neutrosophic sets and called it fuzzy neutrosophic weakly- generalized closed sets in fuzzy neutrosophic topological spaces, we discuss some new properties and examples based of this define concept.

## 1. Basic definitions and terms

### Definition (1.1) [7, 9]:

Let  $X$  be a non-empty fixed set, the fuzzy neutrosophic set (Briefly, FNS),  $\lambda_N$  is an object having the form  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  where the functions  $\mu_{\lambda_N}, \sigma_{\lambda_N}, \nu_{\lambda_N} : X \rightarrow$

\* Corresponding author at: Department of Mathematics, College of Education for Pure Sciences, Tikrit University, Tikrit, IRAQ  
.E-mail address: [nafea\\_v2011@yahoo.com](mailto:nafea_v2011@yahoo.com)

$[0, 1]$ . Denote the degree of membership function (namely  $\mu_{\lambda_N}(x)$ ), the degree of indeterminacy function (namely  $\sigma_{\lambda_N}(x)$ ) and the degree of non-membership (namely  $\nu_{\lambda_N}(x)$ ) respectively, of each set  $\lambda_N$  we have,  $0 \leq \mu_{\lambda_N}(x) + \sigma_{\lambda_N}(x) + \nu_{\lambda_N}(x) \leq 3$ , for each  $x \in X$ .

**Remark (1.2) [9]:**

FNS  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  can be identified to an ordered triple  $\langle X, \mu_{\lambda_N}, \sigma_{\lambda_N}, \nu_{\lambda_N} \rangle$  in  $[0, 1]$  on  $X$ .

**Definition (1.3) [6]:**

Let  $X$  be a non-empty set and the FNSs  $\lambda_N$  and  $\beta_N$  on  $X$  be in the form:

$\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  and,  $\beta_N = \{ \langle x, \mu_{\beta_N}(x), \sigma_{\beta_N}(x), \nu_{\beta_N}(x) \rangle : x \in X \}$  then:

- i.  $\lambda_N \subseteq \beta_N$  iff  $\mu_{\lambda_N}(x) \leq \mu_{\beta_N}(x), \sigma_{\lambda_N}(x) \leq \sigma_{\beta_N}(x)$  and  $\nu_{\lambda_N}(x) \geq \nu_{\beta_N}(x)$  for all  $x \in X$ ,
- ii.  $\lambda_N = \beta_N$  iff  $\lambda_N \subseteq \beta_N$  and  $\beta_N \subseteq \lambda_N$ ,
- iii.  $\underline{1}_N - \lambda_N = \{ \langle x, \nu_{\lambda_N}(x), 1 - \sigma_{\lambda_N}(x), \mu_{\lambda_N}(x) \rangle : x \in X \}$ ,
- iv.  $\lambda_N \cup \beta_N = \{ \langle x, \text{Max}(\mu_{\lambda_N}(x), \mu_{\beta_N}(x)), \text{Max}(\sigma_{\lambda_N}(x), \sigma_{\beta_N}(x)), \text{Min}(\nu_{\lambda_N}(x), \nu_{\beta_N}(x)) \rangle : x \in X \}$ ,
- v.  $\lambda_N \cap \beta_N = \{ \langle x, \text{Min}(\mu_{\lambda_N}(x), \mu_{\beta_N}(x)), \text{Min}(\sigma_{\lambda_N}(x), \sigma_{\beta_N}(x)), \text{Max}(\nu_{\lambda_N}(x), \nu_{\beta_N}(x)) \rangle : x \in X \}$ ,
- vi.  $\underline{0}_N = \langle x, 0, 0, 1 \rangle$  and  $\underline{1}_N = \langle x, 1, 1, 0 \rangle$ .

**Definition (1.4) [9]:**

A fuzzy neutrosophic topology (Briefly, FNT) on a non-empty set  $X$  is a family  $\tau$  of fuzzy neutrosophic subsets in  $X$  satisfying the following axioms:

- i.  $\underline{0}_N, \underline{1}_N \in \tau$ ,
- ii.  $O_{N1} \cap O_{N2} \in \tau$  for any  $O_{N1}, O_{N2} \in \tau$ ,
- iii.  $\cup O_{Ni} \in \tau, \forall \{O_{Ni} : i \in J\} \subseteq \tau$ .

In this case the pair  $(X, \tau)$  is called fuzzy neutrosophic topological space (Briefly, FNTS). The elements of  $\tau$  are called fuzzy neutrosophic open sets (Briefly,  $F_N$ ). The complements of  $F_N$ -open sets in the FNTS  $(X, \tau)$  are called fuzzy neutrosophic closed sets (Briefly,  $F_N$ -closed sets).

**Definition (1.5) [9]:**

Let  $(X, \tau)$  be FNTS and  $\lambda_N = \langle X, \mu_{\lambda_N}, \sigma_{\lambda_N}, \nu_{\lambda_N} \rangle$  be FNS in  $X$ . Then the fuzzy neutrosophic closure of  $\lambda_N$  (Briefly,  $F_NCl$ ) and fuzzy neutrosophic interior of  $\lambda_N$  (Briefly,  $F_NInt$ ) are defined by:

$F_NCl(\lambda_N) = \cap \{C_N : C_N \text{ is } F_N\text{-closed set in } X \text{ and } \lambda_N \subseteq C_N \}$ ,

$F_NInt(\lambda_N) = \cup \{O_N : O_N \text{ is } F_N\text{-open set in } X \text{ and } O_N \subseteq \lambda_N \}$ .

Note that  $F_NCl(\lambda_N)$  be  $F_N$ -closed set and  $F_NInt(\lambda_N)$  be  $F_N$ -open set in  $X$ . Furthermore

- i.  $\lambda_N$  be  $F_N$ -closed set in  $X$  iff  $F_NCl(\lambda_N) = \lambda_N$ ,
- ii.  $\lambda_N$  be  $F_N$ -open set in  $X$  iff  $F_NInt(\lambda_N) = \lambda_N$ .

**Proposition (1.6) [9]:**

Let  $(X, \tau)$  be FNTS and  $\lambda_N, \beta_N$  are FNSs in  $X$ . Then the following properties hold:

- i.  $F_NInt(\lambda_N) \subseteq \lambda_N$  and  $\lambda_N \subseteq F_NCl(\lambda_N)$ ,
- ii.  $\lambda_N \subseteq \beta_N \implies F_NInt(\lambda_N) \subseteq F_NInt(\beta_N)$  and  $\lambda_N \subseteq \beta_N \implies F_NCl(\lambda_N) \subseteq F_NCl(\beta_N)$ ,

- iii.  $\text{FNInt}(\text{FNInt}(\lambda_N)) = \text{FNInt}(\lambda_N)$  and  $\text{FNCl}(\text{FNCl}(\lambda_N)) = \text{FNCl}(\lambda_N)$ ,
- iv.  $\text{FNInt}(\lambda_N \cap \beta_N) = \text{FNInt}(\lambda_N) \cap \text{FNInt}(\beta_N)$  and  $\text{FNCl}(\lambda_N \cup \beta_N) = \text{FNCl}(\lambda_N) \cup \text{FNCl}(\beta_N)$ ,
- v.  $\text{FNInt}(\underline{1}_N) = \underline{1}_N$  and  $\text{FNCl}(\underline{1}_N) = \underline{1}_N$ ,
- vi.  $\text{FNInt}(\underline{0}_N) = \underline{0}_N$  and  $\text{FNCl}(\underline{0}_N) = \underline{0}_N$ .

**Definition (1.7) [8]:**

FNS  $\lambda_N$  in FNTS  $(X, \tau)$  is called:

- i. Fuzzy neutrosophic regular-open set (Briefly, FNR-open)  
 if  $\lambda_N = \text{FNInt}(\text{FNCl}(\lambda_N))$ .
- ii. Fuzzy neutrosophic regular-closed set (Briefly, FNR-closed)  
 if  $\lambda_N = \text{FNCl}(\text{FNInt}(\lambda_N))$ .
- iii. Fuzzy neutrosophic semi-open set (Briefly, FNS-open)  
 if  $\lambda_N \subseteq \text{FNCl}(\text{FNInt}(\lambda_N))$ .
- iv. Fuzzy neutrosophic semi-closed set (Briefly, FNS-closed)  
 if  $\text{FNInt}(\text{FNCl}(\lambda_N)) \subseteq \lambda_N$ .
- v. Fuzzy neutrosophic pre-open set (Briefly, FNP-open)  
 if  $\lambda_N \subseteq \text{FNInt}(\text{FNCl}(\lambda_N))$ .
- vi. Fuzzy neutrosophic pre-closed set (Briefly, FNP-closed)  
 if  $\text{FNCl}(\text{FNInt}(\lambda_N)) \subseteq \lambda_N$ .
- vii. Fuzzy neutrosophic  $\alpha$ -open set (Briefly,  $\text{FN}\alpha$ -open)  
 if  $\lambda_N \subseteq \text{FNInt}(\text{FNCl}(\text{FNInt}(\lambda_N)))$ .
- viii. Fuzzy neutrosophic  $\alpha$ -closed set (Briefly,  $\text{FN}\alpha$ -closed)  
 if  $\text{FNCl}(\text{FNInt}(\text{FNCl}(\lambda_N))) \subseteq \lambda_N$ .

**2. Characterizations and properties of Fuzzy Neutrosophic Weakly-Generalized Closed Sets in Fuzzy Neutrosophic Topological Spaces.**

In this section we introduce and investigate some characterizations and several properties concerning of Fuzzy Neutrosophic Weakly-Generalized Closed Sets in Fuzzy Neutrosophic Topological spaces.

**Definition (2.1) :** Fuzzy neutrosophic sub set  $\lambda_N$  of FNTS  $(X, \tau)$  is called:

- i. Fuzzy neutrosophic-generalized closed set (Briefly, FNGCS) if  $\text{FNCl}(\lambda_N) \subseteq U_N$  whenever,  $\lambda_N \subseteq U_N$  and  $U_N$  be  $F_N$ -open set in  $X$ . And  $\lambda_N$  is said to be fuzzy neutrosophic-generalized open set (Briefly, FNGOS) if the complement  $\underline{1}_N - \lambda_N$  be FNGCS set in  $(X, \tau)$ .
- ii. Fuzzy neutrosophic weakly-closed set (Briefly, FNWCS)  
 if  $\text{FNCl}(\lambda_N) \subseteq U_N$  whenever,  $\lambda_N \subseteq U_N$  and  $U_N$  be FNS-open set in  $X$ . And  $\lambda_N$  is said to be fuzzy neutrosophic weakly-open set (Briefly, FNWOS) if the complement  $\underline{1}_N - \lambda_N$  is FNWCS in  $(X, \tau)$ .
- iii. Fuzzy neutrosophic weakly-generalized closed set (Briefly, FNWGCS) if  $\text{FNCl}(\text{FNInt}(\lambda_N)) \subseteq U_N$  whenever,  $\lambda_N \subseteq U_N$  and  $U_N$  be  $F_N$ -open set in  $X$ . And  $\lambda_N$  is a said to be fuzzy neutrosophic weakly-generalized open set (Briefly, FNWGOS) if the complement  $\underline{1}_N - \lambda_N$  is FNWGCS in  $(X, \tau)$ .

**Theorem (2.2):**

For every FNS, the following statements satisfy:

- i. Every  $F_N$ -closed set is FNGCS.
- ii. Every  $FN\alpha$ -closed set is FNWGCS.
- iii. Every  $F_N$ -closed set is FNWGCS.
- iv. Every FNR-closed set is FNWGCS.
- v. Every FNP-closed set is FNWGCS.

**Proof:**

i. Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be  $F_N$ -closed set in FNTS  $(X, \tau)$ . Then,  $FNCl(\lambda_N) = \lambda_N$ .

Now, let  $\beta_N$  be  $F_N$ -open set such that,  $\lambda_N \subseteq \beta_N$ .

Therefore,  $FNCl(\lambda_N) = \lambda_N \subseteq \beta_N$ .

Hence,  $\lambda_N$  be FNGCS in  $(X, \tau)$ .

ii. Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be  $FN\alpha$ -closed set in FNTS  $(X, \tau)$ . Then,  $FNCl(FNInt(FNCl(\lambda_N))) \subseteq \lambda_N$ .

Now, let  $\beta_N$  be  $F_N$ -open set such that,  $\lambda_N \subseteq \beta_N$ .

Then,  $FNCl(FNInt(\lambda_N)) \subseteq FNCl(FNInt(FNCl(\lambda_N))) \subseteq \lambda_N \subseteq \beta_N$

Therefore,  $FNCl(FNInt(\lambda_N)) \subseteq \beta_N$ .

Hence,  $\lambda_N$  be FNWGCS in  $(X, \tau)$ .

iii. Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be  $F_N$ -closed set in FNTS  $(X, \tau)$ . Then by Definition (1.5) (i),

we get  $\lambda_N = FNCl(\lambda_N) \dots \dots (1)$

By Proposition (1.6) (i) we get,  $FNInt(\lambda_N) \subseteq \lambda_N \dots \dots (2)$

But,  $FNCl(FNInt(\lambda_N)) \subseteq FNCl(\lambda_N)$ .

So by (1),  $FNCl(FNInt(\lambda_N)) \subseteq \lambda_N$

Now, let  $\beta_N$  be  $F_N$ -open set such that,  $\lambda_N \subseteq \beta_N$ .

Then,  $FNCl(FNInt(\lambda_N)) \subseteq \lambda_N \subseteq \beta_N$ .

Therefore,  $FNCl(FNInt(\lambda_N)) \subseteq \beta_N$ .

Hence,  $\lambda_N$  be FNWGCS in  $(X, \tau)$ .

iv. Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be FNR-closed set in FNTS  $(X, \tau)$ . Then,  $FNCl(FNInt(\lambda_N)) = \lambda_N$

Now, let  $\beta_N$  be  $F_N$ -open set such that,  $\lambda_N \subseteq \beta_N$

Then,  $FNCl(FNInt(\lambda_N)) = \lambda_N \subseteq \beta_N$

Therefore,  $FNCl(FNInt(\lambda_N)) \subseteq \beta_N$

Hence,  $\lambda_N$  be FNWGCS in  $(X, \tau)$ .

v. Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be FNP-closed set in FNTS  $(X, \tau)$ . Then,  $FNCl(FNInt(\lambda_N)) \subseteq \lambda_N$

Now, let  $\beta_N$  be  $F_N$ -open set such that,  $\lambda_N \subseteq \beta_N$

Then,  $FNCl(FNInt(\lambda_N)) \subseteq \lambda_N \subseteq \beta_N$

Therefore,  $FNCl(FNInt(\lambda_N)) \subseteq \beta_N$

Hence,  $\lambda_N$  be FNWGCS in  $(X, \tau)$ .

**Remark (2.3) :**

The convers of Theorem (2.2) is not true in general and omit it, it is significant to show it by the following examples:

**Examples (2.4) :**

i. Let  $X = \{a, b\}$  define FNS  $\lambda_N$  in  $X$  as follows:

$$\lambda_N = \langle x, (\frac{a}{0.5}, \frac{b}{0.2}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.5}, \frac{b}{0.7}) \rangle.$$

The family  $\tau = \{0_N, 1_N, \lambda_N\}$  be FNTS.

$$\text{Now if, } \omega_N = \langle x, (\frac{a}{0.9}, \frac{b}{0.3}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.1}, \frac{b}{0.6}) \rangle.$$

And,  $U_N = \underline{1}_N$ , where  $U_N$  be  $F_N$ -open set such that,  $\omega_N \subseteq U_N$ .

Then,  $FNCl(\omega_N) = \bigcap \{C_N : C_N \text{ is } F_N\text{-closed set in } X \text{ and } \omega_N \subseteq C_N\}$

$$= \langle x, (\frac{a}{0.9}, \frac{b}{0.3}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.1}, \frac{b}{0.6}) \rangle \subseteq \langle x, (\frac{a}{1}, \frac{b}{1}), (\frac{a}{1}, \frac{b}{1}), (\frac{a}{0}, \frac{b}{0}) \rangle > \text{ such that,}$$

$$(\frac{a}{0.9}, \frac{b}{0.3}) \leq (\frac{a}{1}, \frac{b}{1}), (\frac{a}{0.5}, \frac{b}{0.5}) \leq (\frac{a}{1}, \frac{b}{1}) \text{ and } (\frac{a}{0.1}, \frac{b}{0.6}) \geq (\frac{a}{0}, \frac{b}{0}) = 1_N.$$

Therefore,  $\text{FNCl}(\omega_N) \subseteq U_N$ .

Hence,  $\omega_N$  is FNGCS but, not  $F_N$ -closed set.

ii. Let  $X=\{a, b\}$  define FNS  $\lambda_N$  in  $X$  as follows:

$$\lambda_N = \langle x, (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.4}, \frac{b}{0.5}) \rangle$$

The family  $\tau = \{0_N, 1_N, \lambda_N\}$  be FNTS,

$$\text{Now if, } \omega_N = \langle x, (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.5}) \rangle.$$

And,  $U_N = \lambda_N$  be  $F_N$ -open set such that,  $\omega_N \subseteq U_N$ .

Then,  $\text{FNInt}(\omega_N) = \cup \{O_N: O_N \text{ is } F_N\text{-open set in } X \text{ and } O_N \subseteq \omega_N\}$

$$= \langle x, (\frac{a}{0}, \frac{b}{0}), (\frac{a}{0}, \frac{b}{0}), (\frac{a}{1}, \frac{b}{1}) \rangle \subseteq \langle x, (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.5}) \rangle.$$

$$\text{Such that, } (\frac{a}{0}, \frac{b}{0}) \leq (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{0}, \frac{b}{0}) \leq (\frac{a}{0.5}, \frac{b}{0.5})$$

$$\text{and } (\frac{a}{1}, \frac{b}{1}) \geq (\frac{a}{0.6}, \frac{b}{0.5}) = 0_N. \text{ And } \text{FNCl}(\text{FNInt}(\omega_N)) = 0_N.$$

Therefore,  $\text{FNCl}(\text{FNInt}(\omega_N)) \subseteq U_N$ .

$$\text{Since, } \langle x, (\frac{a}{0}, \frac{b}{0}), (\frac{a}{0}, \frac{b}{0}), (\frac{a}{1}, \frac{b}{1}) \rangle \subseteq \langle x, (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.4}, \frac{b}{0.5}) \rangle > \text{ such that, } (\frac{a}{0}, \frac{b}{0}) \leq (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0}, \frac{b}{0}) \leq (\frac{a}{0.5}, \frac{b}{0.5}) \text{ and } (\frac{a}{1}, \frac{b}{1}) \geq (\frac{a}{0.4}, \frac{b}{0.5}).$$

Hence,  $\omega_N$  is FNWGCS.

But,  $\text{FNCl}(\omega_N) = \cap \{C_N: C_N \text{ is } F_N\text{-closed set in } X \text{ and } \omega_N \subseteq C\}$

$$= \langle x, (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.5}) \rangle \subseteq \langle x, (\frac{a}{1}, \frac{b}{1}), (\frac{a}{1}, \frac{b}{1}), (\frac{a}{0}, \frac{b}{0}) \rangle > \text{ such that,}$$

$$(\frac{a}{0.5}, \frac{b}{0.4}) \leq (\frac{a}{1}, \frac{b}{1}), (\frac{a}{0.5}, \frac{b}{0.5}) \leq (\frac{a}{1}, \frac{b}{1}) \text{ and } (\frac{a}{0.6}, \frac{b}{0.5}) \geq (\frac{a}{0}, \frac{b}{0}) = 1_N. \text{ So,}$$

$$\text{FNInt}(\text{FNCl}(\omega_N)) = 1_N \text{ and } \text{FNCl}(\text{FNInt}(\text{FNCl}(\omega_N))) = 1_N.$$

Therefore,  $\text{FNCl}(\text{FNInt}(\text{FNCl}(\omega_N))) \not\subseteq \omega_N$ .

$$\text{Since, } \langle x, (\frac{a}{1}, \frac{b}{1}), (\frac{a}{1}, \frac{b}{1}), (\frac{a}{0}, \frac{b}{0}) \rangle \not\subseteq \langle x, (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.5}) \rangle > \text{ such that, } (\frac{a}{1}, \frac{b}{1}) \not\leq (\frac{a}{0.5}, \frac{b}{0.4}), (\frac{a}{1}, \frac{b}{1}) \not\leq (\frac{a}{0.5}, \frac{b}{0.5}) \text{ and } (\frac{a}{0}, \frac{b}{0}) \not\geq (\frac{a}{0.6}, \frac{b}{0.5}).$$

Hence,  $\omega_N$  be not  $FN\alpha$ -closed set.

iii. Take the example which defined in (ii). Then, we can see  $\omega_N$  be FNWGCS but, not  $F_N$ -closed set.

iv. Take again, the example which defined in (ii). Then,  $\omega_N$  be FNWGCS but not  $F_N$ -closed set.

v. Take, the example which defined in (i). Then,  $\text{FNInt}(\omega_N) = \lambda_N$  and  $\text{FNCl}(\text{FNInt}(\omega_N)) = 1_N \cdot \lambda_N$ .

Therefore,  $\text{FNCl}(\text{FNInt}(\omega_N)) \subseteq U_N$ .

Hence,  $\omega_N$  is FNWGCS.

But,  $\text{FNCl}(\text{FNInt}(\omega_N)) \not\subseteq \omega_N$ .

Hence,  $\omega_N$  is not  $F_N$ -closed set.

### Remark (2.5) :

The relation between FNS-closed sets and FNWGCSs is independent, it is important to show it by the following examples:

### Example (2.6) :

(1) Let  $X=\{a, b\}$  define FNS  $\lambda_N$  in  $X$  as follows:

$$\lambda_N = \langle x, (\frac{a}{0.3}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.7}) \rangle.$$

The family  $\tau = \{0_N, 1_N, \lambda_N\}$  be FNTS.

$$\text{Now if, } \omega_N = \langle x, (\frac{a}{0.3}, \frac{b}{0.4}), (\frac{a}{0.5}, \frac{b}{0.5}), (\frac{a}{0.6}, \frac{b}{0.7}) \rangle,$$

And,  $U_N = \lambda_N$  where  $U_N$  be  $F_N$ -open set such that,  $\omega_N \subseteq U_N$ .

Then,  $FNCI(\omega_N) = \underline{1}_N - \lambda_N$  and  $FNInt(FNCI(\omega_N)) = \lambda_N$ .

Therefore,  $FNInt(FNCI(\omega_N)) \subseteq \omega_N$ .

Hence,  $\omega_N$  is FNS-closed set.

But,  $FNInt(\omega_N) = \lambda_N$  and  $FNCI(FNInt(\omega_N)) = \underline{1}_N - \lambda_N$ .

Therefore,  $FNCI(FNInt(\omega_N)) \not\subseteq U_N$ .

Hence,  $\omega_N$  is not FNWGCS.

(2) Take Example (2.4) (v) then,  $\omega_N$  is FNWGCS.

But,  $FNCI(\omega_N) = \underline{1}_N$  and  $FNInt(FNCI(\omega_N)) = \underline{1}_N$ .

Therefore,  $FNInt(FNCI(\omega_N)) \not\subseteq \omega_N$ .

Hence,  $\omega_N$  is not FNS-closed set.

**Proposition (2.7) :**

Let  $\lambda_N$  be  $F_N$ -closed set in  $(X, \tau)$  such that,  $FNInt(\lambda_N) \subseteq \beta_N \subseteq \lambda_N$ . Then,  $\beta_N$  is FNWGCS on FNTS  $(X, \tau)$ .

**Proof:** Let  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be FNS in FNTS  $(X, \tau)$  such that,  $FNInt(\lambda_N) \subseteq \beta_N \subseteq \lambda_N$ .

So, there exists  $F_N$ -closed set  $\eta_N$  such that,  $\eta_N (FNInt(\lambda_N)) \subseteq \beta_N \subseteq \lambda_N \subseteq \eta_N$ .

Then,  $\beta_N \subseteq \eta_N$  and also  $FNInt(\beta_N) \subseteq \beta_N \subseteq \eta_N$ .

Thus,  $FNCI(FNInt(\beta_N)) \subseteq \beta_N$ .

Now, let  $\Psi_N$  be  $F_N$ -open set such that,  $\beta_N \subseteq \Psi_N$ .

Then,  $FNCI(FNInt(\beta_N)) \subseteq \beta_N \subseteq \Psi_N$ .

Therefore,  $FNCI(FNInt(\beta_N)) \subseteq \Psi_N$ .

Hence,  $\beta_N$  is FNWGCS in  $(X, \tau)$ .

**Theorem (2.8) :**

Let  $(X, \tau)$  be FNTS, then the intersection of two FNWGCSs is also FNWGCS.

**Proof:** Let  $\lambda_N$  and  $\beta_N$  are FNP-closed sets on FNTS  $(X, \tau)$ .

Then,  $FNCI(FNInt(\lambda_N)) \subseteq \lambda_N$  and  $FNCI(FNInt(\beta_N)) \subseteq \beta_N$ .

Consider  $\lambda_N \cap \beta_N \supseteq FNCI(FNInt(\lambda_N)) \cap FNCI(FNInt(\beta_N))$

$$\supseteq FNCI(FNInt(\lambda_N) \cap FNInt(\beta_N))$$

$$\supseteq FNCI(FNInt(\lambda_N \cap \beta_N))$$

This means  $FNCI(FNInt(\lambda_N \cap \beta_N)) \subseteq \lambda_N \cap \beta_N$ .

Now, let  $\eta_N$  be  $F_N$ -open set such that,  $\lambda_N \cap \beta_N \subseteq \eta_N$

Then,  $FNCI(FNInt(\lambda_N \cap \beta_N)) \subseteq \lambda_N \cap \beta_N \subseteq \eta_N$ .

Therefore,  $FNCI(FNInt(\lambda_N \cap \beta_N)) \subseteq \eta_N$ .

Hence,  $\lambda_N \cap \beta_N$  be FNWGCS in  $(X, \tau)$ .

**Remark (2.9) :**

The union of any FNWGCSs is not necessary to be FNWGCS see the following example:

**Example (2.10) :**

Let  $X=\{x\}$  define FNSs  $\lambda_N$  and  $\beta_N$  in  $X$  as follows:

$\lambda_N =\{ \langle x, 0.5, 0.6, 0.7 \rangle : x \in X \}$  and  $\beta_N =\{ \langle x, 0.6, 0.7, 0.5 \rangle : x \in X \}$ .

The family  $\tau = \{ \underline{0}_N, \underline{1}_N, \lambda_N, \beta_N \}$  be FNTS.

Now if,  $\omega_{N1} = \{ \langle x, 0.6, 0.5, 0.5 \rangle : x \in X \}$ ,  $\omega_{N2} = \{ \langle x, 0.6, 0.7, 0.8 \rangle : x \in X \}$  and

$U_N =\{ \langle x, 0.6, 0.7, 0.5 \rangle : x \in X \}$  where,  $U_N$  be  $F_N$ -open set such that,  $\omega_{N1} \subseteq U_N$  and  $\omega_{N2} \subseteq U_N$ .

Then,  $FNInt(\omega_{N1}) = \underline{0}_N$  and  $FNCl(FNInt(\omega_{N1})) = \underline{0}_N$

Therefore,  $FNcl(FNint(\omega_{N1})) \subseteq U_N$ . Hence,  $\omega_{N1}$  be FNWGCS.

And,  $FNint(\omega_{N2}) = \underline{0}_N$  and  $FNcl(FNint(\omega_{N2})) = \underline{0}_N$ .

Therefore,  $FNCl(FNInt(\omega_{N2})) \subseteq U_N$ .

Hence,  $\omega_{N2}$  be FNWGCS.

So,  $\omega_{N1} \cup \omega_{N2} =\{ \langle x, 0.6, 0.7, 0.5 \rangle : x \in X \}$ .

But,  $FNInt(\omega_{N1} \cup \omega_{N2} ) =\{ \langle x, 0.6, 0.7, 0.5 \rangle : x \in X \}$  and  $FNCl(FNInt(\omega_{N1} \cup \omega_{N2})) = \underline{1}_N$

Therefore,  $FNCl(FNInt ( \omega_{N1} \cup \omega_{N2})) \not\subseteq U_N$

Hence,  $\omega_{N1} \cup \omega_{N2}$  is not FNWGCS.

**Definition (2.11) :**

Let  $(X, \tau)$  be FNTS and  $\lambda_N = \{ \langle x, \mu_{\lambda_N}(x), \sigma_{\lambda_N}(x), \nu_{\lambda_N}(x) \rangle : x \in X \}$  be FNS. Then, the fuzzy neutrosophic is weakly generalized closure of  $\lambda_N$  (Briefly, FNWGCl) and fuzzy neutrosophic weakly generalized interior of  $\lambda_N$  (Briefly, FNWGInt) are defined by:

- i.  $FNWGCl(\lambda_N) = \cap \{ \beta_N : \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \}$
- ii.  $FNWGInt(\lambda_N) = \cup \{ \beta_N : \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \}$ ,

**Proposition (2.12):**

Let  $(X, \tau)$  be FNTS and  $\lambda_N, \beta_N$  are FNSs in  $X$ . Then the following properties hold:

- i.  $FNWGCl(0_N) = \underline{0}_N$  and  $FNWGCl(1_N) = \underline{1}_N$ ,
- ii.  $\lambda_N \subseteq FNWGCl(\lambda_N)$ ,
- iii. If  $\lambda_N \subseteq \beta_N$ , then  $FNWGCl(\lambda_N) \subseteq FNWGCl(\beta_N)$ ,
- iv.  $\lambda_N$  is FNWGCS iff  $\lambda_N = FNWGCl(\lambda_N)$
- v.  $FNWGCl(FNWGCl(\lambda_N)) = FNWGCl(\lambda_N)$ .

**Proof:**

- i. By Definition (2.11) (i) it is important to focus on:

$FNWGCl(0_N) = \cap \{ \beta_N : \beta_N \text{ is FNWGCS in } X \text{ and } 0_N \subseteq \beta_N \} = \underline{0}_N$ ,

And,  $FNWGCl(1_N) = \cap \{ \beta_N : \beta_N \text{ is FNWGCS in } X \text{ and } 1_N \subseteq \beta_N \} = \underline{1}_N$ .

- ii.  $\lambda_N \subseteq \cap \{ \beta_N : \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \} = FNWGCl(\lambda_N)$ .

iii. Suppose that  $\lambda_N \subseteq \beta_N$  then,  $\cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \}$

$$\subseteq \cap \{ \eta_N: \eta_N \text{ is FNWGCS in } X \text{ and } \beta_N \subseteq \eta_N \}$$

Therefore,  $\text{FNWGCI}(\lambda_N) \subseteq \text{FNWGCI}(\beta_N)$ .

iv.  $\Rightarrow$  If,  $\lambda_N$  is FNWGCS then,

$$\text{FNWGCI}(\lambda_N) = \cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \} \dots \dots (1)$$

$$\text{And by (ii), } \lambda_N \subseteq \text{FNWGCI}(\lambda_N) \dots \dots (2)$$

But  $\lambda_N$  is necessarily to be the smallest set.

Thus,  $\lambda_N = \cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \}$ ,

$$\text{Therefore, } \lambda_N = \text{FNWGCI}(\lambda_N)$$

$\Leftarrow$  Let  $\lambda_N = \text{FNWGCI}(\lambda_N) = \cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \}$  and by using Definition 2.11 (i), we get  $\lambda_N$  is FNWGCS in X.

v. Since,  $\lambda_N = \text{FNWGCI}(\lambda_N)$  so we get,  $\text{FNWGCI}(\lambda_N) = \text{FNWGCI}(\text{FNWGCI}(\lambda_N))$ .

**Proposition (2.13):**

Let  $(X, \tau)$  be FNTS and  $\lambda_N, \beta_N$  are FNSs in X. Then the following properties hold:

i.  $\text{FNWGInt}(0_N) = \underline{0}_N$  and  $\text{FNWGInt}(1_N) = \underline{1}_N$ ,

ii.  $\text{FNWGInt}(\lambda_N) \subseteq \lambda_N$ ,

iii. If  $\lambda_N \subseteq \beta_N$ , then  $\text{FNWGInt}(\lambda_N) \subseteq \text{FNWGInt}(\beta_N)$ ,

iv.  $\lambda_N$  is a FNWGOS iff  $\lambda_N = \text{FNWGInt}(\lambda_N)$ ,

v.  $\text{FNWGInt}(\lambda_N) = \text{FNWGInt}(\text{FNWGInt}(\lambda_N))$ .

**Proof:**

i. By Definition (2.11) (ii) we have

$$\text{FNWGInt}(0_N) = \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq 0_N \} = \underline{0}_N$$

$$\text{And, } \text{FNWGInt}(1_N) = \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq 1_N \} = \underline{1}_N$$

ii. Follows from Definition (2.11) (ii).

iii.  $\text{FNWGInt}(\lambda_N) = \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \}$ .

Since,  $\lambda_N \subseteq \beta_N$  then,  $\cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \}$

$$\subseteq \cup \{ \eta_N: \eta_N \text{ is FNWGOS in } X \text{ and } \eta_N \subseteq \beta_N \}$$

$$\text{Therefore, } \text{FNWGInt}(\lambda_N) \subseteq \text{FNWGInt}(\beta_N)$$

iv.  $\Rightarrow$  Omit it, there must be a proof that  $\text{FNWGInt}(\lambda_N) \subseteq \lambda_N$  and  $\lambda_N \subseteq \text{FNWGInt}(\lambda_N)$ . Suppose that  $\lambda_N$  is FNWGOS in X.

Then,  $\text{FNWGInt}(\lambda_N) = \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \}$  by using ii we get,  $\text{FNWGInt}(\lambda_N) \subseteq \lambda_N \dots \dots (1)$

Now to proof,  $\lambda_N \subseteq \text{FNWGInt}(\lambda_N)$ , we have, for all  $\lambda_N \subseteq \lambda_N$ ,

The  $\text{FNWGInt}(\lambda_N) \subseteq \lambda_N$ . So, we get

$$\lambda_N \subseteq \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \} = \text{FNWGInt}(\lambda_N) \dots \dots (2)$$

From (1) and (2) we have,  $\lambda_N = \text{FNWGInt}(\lambda_N)$ .

$\Leftarrow$  Suppose that  $\lambda_N = \text{FNWGInt}(\lambda_N)$  and by using Definition 2.11 (ii), we

get  $\lambda_N$  is a FNWGOS in X.



v. Since,  $\lambda_N = \text{FNWGInt}(\lambda_N)$  by (iv) so we get,

$$\text{FNWGInt}(\lambda_N) = \text{FNWGInt} (\text{FNWGInt} (\lambda_N)).$$

**Theorem (2.14):**

Let  $(X, \tau)$  be FNTS. Then for every fuzzy neutrosophic subsets  $\lambda_N$  of X, we have:

- i.  $\underline{1}_N - \text{FNWGInt} (\lambda_N) = \text{FNWGCI} (\underline{1}_N - \lambda_N),$
- ii.  $\underline{1}_N - \text{FNWGCI} (\lambda_N) = \text{FNWGInt} (\underline{1}_N - \lambda_N).$

**Proof:**

i.  $\text{FNWGInt}(\lambda_N) = \cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \}$ , by the complement, we get  $\underline{1}_N - \text{FNWGInt} (\lambda_N) = \underline{1}_N - (\cup \{ \beta_N: \beta_N \text{ is FNWGOS in } X \text{ and } \beta_N \subseteq \lambda_N \})$ .

So,  $\underline{1}_N - \text{FNWGInt}(\lambda_N) = \cap \{ (1_N - \beta_N): (1_N - \beta_N) \text{ is FNWGCS in } X \text{ and } (\underline{1}_N - \lambda_N) \subseteq (\underline{1}_N - \beta_N) \}$ .

Now, replacing  $(\underline{1}_N - \beta_N)$  by  $\eta_N$  we get,

$$\begin{aligned} \underline{1}_N - \text{FNWGInt}(\lambda_N) &= \cap \{ \eta_N: \eta_N \text{ is FNWGCS in } X \text{ and } (\underline{1}_N - \lambda_N) \subseteq \eta_N \} \\ &= \text{FNWGCI}(\underline{1}_N - \lambda_N). \end{aligned}$$

ii.  $\text{FNWGCI}(\lambda_N) = \cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \}$ , by the complement, we get  $\underline{1}_N - \text{FNWGCI}(\lambda_N) = \underline{1}_N - (\cap \{ \beta_N: \beta_N \text{ is FNWGCS in } X \text{ and } \lambda_N \subseteq \beta_N \})$

So,  $\underline{1}_N - \text{FNWGCI} (\lambda_N) = \cup \{ (\underline{1}_N - \beta_N) : (\underline{1}_N - \beta_N) \text{ is FNWGOS in } X \text{ and } (\underline{1}_N - \beta_N) \subseteq (\underline{1}_N - \lambda_N) \}$ .

Again replacing  $(\underline{1}_N - \beta_N)$  by  $\eta_N$  we get,

$$\begin{aligned} \underline{1}_N - \text{FNWGCI}(\lambda_N) &= \cup \{ \eta_N: \eta_N \text{ is FNWGOS in } X \text{ and } \eta_N \subseteq (\underline{1}_N - \lambda_N) \} \\ &= \text{FNWGInt}(\underline{1}_N - \lambda_N). \end{aligned}$$

**Theorem (2.15) :**

If  $(X, \tau)$  be FNTS, then for every fuzzy neutrosophic subsets  $\lambda_N$  and  $\beta_N$  of X we have

- i.  $\lambda_N \cup \text{FNWGCI} (\text{FNWGInt} (\lambda_N)) \subseteq \text{FNWGCI} (\lambda_N),$
- ii.  $\text{FNWGInt} (\lambda_N) \subseteq \lambda_N \cap \text{FNWGInt} (\text{FNWGCI} (\lambda_N)).$

**Proof:**

i. Since, by Proposition (2.12) (ii)  $\lambda_N \subseteq \text{FNWGCI}(\lambda_N)$ .....(1)

and, by Proposition (2.13) (ii) we have,  $\text{FNWGInt} (\lambda_N) \subseteq \lambda_N$

Then,  $\text{FNWGCI} (\text{FNWGInt} (\lambda_N)) \subseteq \text{FNWGCI} (\lambda_N)$ .....(2)

So, from (1) and (2) we get,

$$\lambda_N \cup \text{FNWGCI} (\text{FNWGInt}(\lambda_N)) \subseteq \text{FNWGCI}(\lambda_N).$$

ii. Since, by Proposition (2.13) (ii) we have,  $\text{FNWGInt} (\lambda_N) \subseteq \lambda_N$ .....(\*) and,


by Proposition (2.12) (ii) we have,  $\lambda_N \subseteq \text{FNWGCI} (\lambda_N)$


Then,  $\text{FNWGInt}(\lambda_N) \subseteq \text{FNWGInt} (\text{FNWGCI} (\lambda_N))$ .....(\*\*)

So, from (\*) and (\*\*) we get,

$$\text{FNWGInt} (\lambda_N) \subseteq \lambda_N \cap \text{FNWGInt} (\text{FNWGCI} (\lambda_N)).$$

**Remark (2.16):** The relationship between different sets in FNTS see the next Figure-1 and the convers is not true in general.





**(Figure-1) The relationship between different sets of Fuzzy Neutrosophic Topological Space Conclusion**

In this paper, we defined new class of sets, is the fuzzy Neutrosophic weakly- generalized closed sets, then we proved some theorems related to this definition. We introduced defined for the new class of sets by fuzzy Neutrosophic sets and called it the fuzzy Neutrosophic weakly-generalized closed sets in fuzzy Neutrosophic topological spaces, we discuss some new properties, theorems and examples based of this define concept.

#### References:

[1] L. A. Zadeh. (1965). *Fuzzy Sets*. Inform and Control, Vol.8, 338:353.

[2] K. Atanassov and S. Stoeva. (August 1983). *Intuitionistic Fuzzy Sets*. in: Polish Syrup. On Interval & Fuzzy Mathematics, Poznan, 23:26.

[3] K. Atanassov. (1986). *Intuitionistic Fuzzy Sets*, Fuzzy Sets and Systems, Vol. 20, 87:96.

[4] K. Atanassov. (1988). *Review and new results on Intuitionistic Fuzzy Sets*, Preprint IM- MFAIS, Sofia. 1:88.

[5] F. Smaradache. (2010). *Neutrosophic Set : A Generalization of Intuitionistic Fuzzy Set*. Journal of Defense Resources Management. Vol. 1, 1:10.

[6] A. A. Salama and S. A. Alblowi. (2012). *Neutrosophic Set and Neutrosophic Topological Spaces*. IOSR Journal of Mathematics, Vol. 3, Iss. 4. 31:35.

[7] I.Arockiarani and J.Martina Jency. (2014). *More on Fuzzy Neutrosophic Sets and Fuzzy Neutrosophic Topological Spaces*. IJIRS, Vol. 3. 642:652.

[8] I. Arokiarani, R. Dhavaseelan, S. Jafari and M. Parimala. (2017). *On Some Notions and Functions in Neutrosophic Topological Spaces*. Neutrosophic Sets and System. Vol. 16, 16:19.

[9] Y.Veeresswari , *An Introduction To Fuzzy Neutrosophic Topological Spaces*, IJMA, Vol.8(3), (2017). 144-149.

## المجموعات المغلقة المعممة - الضعيفة النايتروسوفيك المضببة في الفضاءات التوبولوجية النايتروسوفيك المضببة

فاطمة محمود محمد, انس عباس حجاب, شيماء فائق مطر

قسم الرياضيات - كلية التربية للعلوم الصرفة - جامعة تكريت

[nafea\\_y2011@yahoo.com](mailto:nafea_y2011@yahoo.com)

#### الخلاصة:

في هذا البحث، قمنا بتعريف فئة جديدة من المجموعات تم تسميتها بالمجموعات المغلقة المعممة - الضعيفة النايتروسوفيك المضببة، ثم أثبتنا بعض النظريات المتعلقة بهذا التعريف. بعد ذلك درسنا بعض العلاقات بين المجموعات المغلقة المعممة - الضعيفة النايتروسوفيك المضببة من جهة والمجموعات المغلقة  $\alpha$  نيوتروسوفيك المضببة، المجموعات المغلقة النايتروسوفيك المضببة، المجموعات المغلقة المنتظمة نيوتروسوفيك المضببة، المجموعات المغلقة القبلية نيوتروسوفيك المضببة والمجموعات شبه المغلقة نيوتروسوفيك المضببة من جهة أخرى في الفضاءات التوبولوجية نيوتروسوفيك المضببة مع بعض الخصائص.